

RINGEL'S GENERALIZED EARTH-MOON PROBLEM

JEREMY ALEXANDER

1. INTRODUCTION

A graph G has surface thickness n (S -thickness n) with respect to the surface S , if G can be decomposed into n and no fewer subgraphs by making n copies of the vertex set of G and then assigning each edge of G to one of the n copies so that n graphs result; each resulting subgraph of G must be S embeddable. The chromatic number of a graph G , denoted $\chi(G)$, is simply the fewest number of colors needed to color the vertex set of G such that no two adjacent vertices receive the same color. Define $\chi_n(S)$ to be the chromatic number of the surface S , that is, the minimum number of colors needed to properly color all S -thickness n graphs. Let $E(S)$ denote the Euler characteristic of the surface S . Let S_k denote the orientable surface of genus k and N_k the nonorientable surface obtained by adding k crosscaps to the sphere; S_0 denotes the two-dimensional sphere. It is known that $E(S_k) = 2 - 2k$ and $E(N_k) = 2 - k$.

In 1959 [1, p.233], Ringel asked: What is the chromatic number of the sphere for graphs of thickness two? The bounds are known to be: $9 \leq \chi_2(S_0) \leq 12$. See [2] for a delightful exposition of the spherical case. In [1], Ringel and Jackson ask: What is $\chi_n(S)$ for any surface S ? There [p.241] the following upper bound is given for any surface S , except the sphere:

$$\chi_n(S) \leq \left\lfloor (1 + 6n + \sqrt{(1 + 6n)^2 - 24nE(S)}) / 2 \right\rfloor \quad (1)$$

The main results in this note are three elementary arguments that establish $\chi_n(S)$ for two new surfaces: N_3 , also known as Dyck's surface, and the Moebius band which will be denoted by M . The arguments are based on previous results appearing in [1]. The results are not especially surprising, but they do provide good lower bounds for $\chi_n(N_{2k+1})$ once the value of $\chi_n(S_k)$ is known for $k > 0$; in the case $k = 1$, the exact value of $\chi_n(N_3)$ is obtained.

2. MAIN RESULTS

Lemma 1. $\left\lfloor (1+6n + \sqrt{(1+6n)^2 - 24nE(N_3)}) / 2 \right\rfloor \leq 6n+1$ for all $n \geq 2$.

Proof. Suppose for contradiction that the claim is false.

Then since $E(N_3) = -1$, there exists an integer $n \geq 2$ such that:

$$\left\lfloor (1+6n + \sqrt{(1+6n)^2 + 24n}) / 2 \right\rfloor > 6n+1 \rightarrow$$

$$\left\lfloor (1+6n + \sqrt{(1+6n)^2 + 24n}) / 2 \right\rfloor \geq 6n+2 \rightarrow$$

$$1+6n + \sqrt{(1+6n)^2 + 24n} \geq 12n+4 \rightarrow$$

$$\sqrt{(1+6n)^2 + 24n} \geq 6n+3 \rightarrow$$

$$(1+6n)^2 + 24n \geq (6n+3)^2 \rightarrow$$

$1 \geq 9$. This is an obvious contradiction, hence, the original assumption was false and the lemma is true. $\square$


Theorem 1. $\chi_n(N_3) = 6n+1$ for all $n \geq 2$.

Proof. In [1, p.238], it is shown that $\chi_n(S_1) = 6n+1$ for all $n \geq 2$. Let G be any graph embedded on S_1 . Choose any face of S_1 created by the embedding of G and add a crosscap, what can be obtained is an embedding of G on the surface N_3 since N_3 is homeomorphic to the surface $S_1 + \text{crosscap}$ and homeomorphism preserves graph isomorphism as well as graph embedding. Combining this with the previous lemma gives the desired result. $\square$

For a complete classification of surfaces see Conway's ZIP proof in [3].

Theorem 2. $\chi_n(M) = 6n$ for all $n \geq 2$.

Proof. M is homeomorphic to N_1 minus an open disk, to see this look at the figure below; a square minus the region X (an open disk) with opposite ends identified. Notice that coupled ends have opposite orientation and so a half-twist is needed to join each pair correctly. When opposite ends are joined a Moebius band results. Now, fill in the two missing regions denoted by X . If sides of the resulting square receive the same identifications induced by the labeling of the figure below, N_1 , the projective plane, is obtained. Now, given any graph G embedded on N_1 , select any face induced by the embedding and delete it, what results is an embedding of G on a Moebius band. Inductively it follows that $\chi_n(M) \geq 6n$ for all $n \geq 2$ since in [1, p.240] it is proven that $\chi_n(N_1) = 6n$ for all $n \geq 2$. Now, inequality (1) gives an upper bound of $\chi_n(M) \leq 6n+1$ for all $n \geq 2$ since $E(M) = 0$. To show that this upper bound cannot be obtained, the case $\chi_2(M)$ is shown, from which the general result easily follows. Suppose for contradiction that $\chi_2(M) = 13$. Then there exists a graph G of M -thickness two such that $\chi(G) = 13$. Since the square minus an open disk appearing in the picture below is homeomorphic to M , G biembeds on this version of a Moebius band. Now fill in the deleted disk labeled X in the figure below to obtain a biembedding of the graph G on N_1 . This implies that $\chi_2(N_1) = 13$ which is a contradiction since $\chi_2(N_1) = 12$. $\square$



Hence, Ringel's generalized earth-moon problem has been solved for two new surfaces M and N_3 , also known as the Moebius band and Dyck's surface, respectively. The following list summarizes known results for Ringel's generalized earth-moon problem:

For all $n \geq 2$,

$$\chi_n(S_2) = 6n + 2 \quad (\text{see [5]})$$

$$\chi_n(S_1) = 6n + 1 \quad (\text{see [1], [5]})$$

$$\chi_n(N_1) = 6n \quad (\text{see [1], [5]})$$

$$\chi_n(M) = 6n$$

$$\chi_n(N_3) = 6n + 1.$$

Until very recently the N_2 -thickness of K_{13} was not known, however, it appears that Thom Sulanke has determined it is three using an exhaustive computer search [4]. The N_2 -thickness of K_{13} has been a long standing obstruction to extending Beineke's results appearing in [5] for the Klein Bottle thickness of K_n . A similar obstruction for determining the S_3 -thickness of K_n is the S_3 -thickness of K_{16} [5, p.996]; Sulanke has determined that $\chi_2(S_3) = 16$, although this result has not been published. Through personal communication Thom has also told me that he has been able to determine $\chi_n(N_4)$ and $\chi_n(N_5)$. He also pointed out to me that Lemma 1, appearing here, generalizes for all integers $n > 0$ and $E(S) < 0$. Using the exact same arguments given in the proof of Lemma 1, one can show that the left side of inequality (1) is $\leq 6n - E(S)$ for $n > 0$ and $E(S) < 0$. Naturally, the next two surfaces to consider are S_3 and N_2 . This motivates the following problem:

Problem 1. Determine $\chi_n(S_3)$ and $\chi_n(N_2)$.

Acknowledgement 1. I would like to thank Yo'av Rieck and Chaim Goodman-Strauss for constructive criticism.

3. REFERENCES

- [1] B. Jackson, G. Ringel, Variations on Ringel's earth-moon problem, *Discrete Mathematics* **211** (2000), 233 - 242.
- [2] J. P. Hutchinson, Coloring ordinary maps, maps of empires, and maps of the moon, *Math. Magazine* (4) **66** (1993), 211 - 226.
- [3] G. Francis, J. Weeks, Conway's ZIP proof, *Amer. Math. Monthly* **106** (1999), 393-399.
- [4] T. Sulanke, Biembeddings of K_{13} ,
<http://needmore.physics.indiana.edu/~tsulanke/graphs/biembed/biembed.pdf>
- [5] L. W. Beineke, Minimal decompositions of complete graphs into subgraphs with embeddability properties, *Canad. J. Math* **21** (1969), 992-1000.

MATHEMATICS DEPARTMENT SCEN 301, UNIVERSITY OF ARKANSAS, FAYETTEVILLE, AR 72701

E-mail address: `jralexa@uark.edu`

URL: `comp.uark.edu/~jralexa`